



Seasonal variation of protein composition of *Rastrelliger kanagartha* (Cuvier) from within PFZ and outside PFZ at murud fish landing centre on the coast of raigad district at Maharashtra state

Budharatna Bhaware¹, Digvijay Lawate²

¹ Department of Zoology, G.M. Vedak College of Science, Tala-Raigad, Maharashtra, India

² Department of Zoology, ACS College, Lanja-Ratnagiri, Maharashtra, India

Abstract

The protein content in the muscles, liver and gonad of *Rastrelliger kanagartha* from Murud fish landing centre at Raigad district coast was studied from Within Potential Fishing Zone (PFZ) and Outside PFZ. The protein study analyzed that winter and summer season. The protein content was increased in male muscle than female muscle, gonad and liver respectively within PFZ than outside PFZ. The protein contents in the body muscle, liver and gonad of *Rastrelliger kanagartha* (Cuvier) were investigated. The compositions varied seasonally in relation to reproductive cycle of the fish. The study also showed that protein was high in the muscles and gonads and in liver from within PFZ than outside PFZ in female and male.

Keywords: Protein content, *Rastrelliger Kanagartha*, within PFZ and outside PFZ, murud fish landing centre

Introduction

Satellite based Sea Surface Temperature (SST) images are being used as an input for locating potential fishing zone (PFZ) of productivity and hence fish availability for commercial fishing operations (Pillai, 2005) [1]. The PFZ advisories pertaining to these coastal States are being generated using the satellite data sets and the SST and chlorophyll-a were derived from NOAA AVHRR, IRS-P4 OCM and MODIS AQUA. Fish catch data pertaining to the PFZ advisories were collected from both the boats operating within and outside PFZ. Fishes are rich source of animal protein for the population. The nutritive value of fish is recognized all over the world. Fishes is very important for rich source. It is therefore obvious that an understanding of the chemical constitution and nutritional quality of fish is important in its use as food of human. Reconsideration of the literature shows that biochemical composition of different body parts of fish has not received adequate attention. Fish is a major source of food for human nutrition providing an important amount of dietary protein diet in many countries. Fish flesh is easily digestible because it contains long muscle fibers. Furthermore, it has been linked to health benefits, such as the prevention of cardiovascular diseases and some types of cancer, including colon, breast and prostate (Rose and Connoll, 1993; Marchioli, 2001, 2002; Sidhu, 2003) [2, 3, 4, 5].

Fish is a good source of many important nutrients such as proteins. Fish has long been a popular meal of people living around the Konkan coast and has been a major source of food for people living this region. The biochemical contents of fishes are subject to marked seasonal Potential Fishing Zone and outside PFZ changes, which have been attributed to factors such as maturation, spawning, age, growth and feeding. The mackerel fish, *Rastrelliger kanagartha* is abundant in the Ratnagiri and Raigad district coasts, at Maharashtra State and forms subsistent fishery. This fish is one of the favorite items of the diet of local people. The paper is an attempt to evaluate the variations in proteins in different body tissues like muscle, liver and gonad in

relation to within PFZ and outside PFZ in winter and summer season from males & females' fish.

Materials and Methods

The PFZ advisories are generated by INCOIS thrice in a week (Monday, Wednesday and Friday), in non-monsoon and non-banned period and cloud free days, there were used by the fishermen along Ratnagiri district coasts. Depending on the location of Potential Fishing Zone (PFZ), i.e. the advisories provided by personal contacts, fax or telephone messages and on Digital Display Boards installed at major landing centres. The boats were selected for the within and outside PFZ region in the Murud fish landing centre at Raigad districts. The PFZ boats and outside PFZ were returned to fishing at Murud FLC and collected the feedback from the users, fishing by the purse-seine and trawl. The pelagic fishes such as mackerel *R. kanagartha* caught by purse-seine within PFZ and outside PFZ region. The PFZ vessels using PFZ–Mission forecast and outside PFZ vessels not using the PFZ forecast. The mackerel samples of *R. Kanagartha* were collected from fishermen catches of within and outside PFZ region of Murud. After bringing them to the research laboratory the total length, body weight, sex and stage of maturity were determined based on findings (Nikolsky, 1963) [6]. The fishes washed thoroughly and the surface moisture was removed by blotting paper. The muscular, hepatic and gonadal tissues were dissected, weight to nearest 1 mg using digital electronic balance. The wet weight tissues were taken for further protein analysis. The optical density of the color developed for protein, glycogen and lipid was measured using spectrophotometer. The total protein estimation was made employing the method by (Lowry *et al.* 1951) [7], Glycogen by De-zwaan and Zandee (1972) [8], and Lipid estimation by Barnes and Black stock (1973) [9]. Results were expressed as mg/100mg wet weight tissues, biochemical constituents were correlated within and outside PFZ region. The standard deviations expressed were arithmetic means, S. D. and statistical analysis.

Results

The study showed that the analysis of protein constituent from muscle, liver and gonad in the within and outside PFZ

during winter and summer season. The mean values for each area were calculated and the results were expressed in mg of protein/100mg of wet wt. Tissues

Table 1: Protein (mg/100mg wet wt.) content in male *R. kanagurta* from Murud

Landing centre	Season	Fishing zone	Tissue		
			Muscle	Liver	Gonad
Murud	Winter	Within PFZ	20.04 ± 0.30	12.32 ± 1.04	20.40 ± 1.05 * °°
		Outside PFZ	18.08 ± 0.34	13.20 ± 0.05	19.40 ± 0.54 * °°
	Summer	Within PFZ	21.24 ± 0.45	11.33 ± 0.44	20.01 ± 1.24 *** °°
		Outside PFZ	19.06 ± 0.66	10.56 ± 0.51	18.20 ± 0.64 °°

•=P<0.001, ••= P<0.01, •••=P<0.05 comparison was made muscle to liver
 * =P<0.001, **= P<0.01, ***=P<0.05 comparison was made liver to gonad
 ° =P<0.001, °°= P<0.01, °°° =P<0.05 comparison was made muscle to gonad




Fig 1: Protein (mg/100mg wet wt.) content in male *R. kanagurta* from Murud

Protein

Murud Fish landing centre in male

In Table. 1. & Fig. 1. Shows that males protein content from muscle was 20.04 ± 0.30 and liver 12.32 ± 1.04 while, in testis it was higher 20.40 ± 1.05 from within PFZ and in outside PFZ males muscle protein content was 18.08 ± 0.34 and in liver 13.20 ± 0.05 while in testis it was protein content was 19.40 ± 0.54 during winter season. The protein content in males' muscle was 21.24 ± 0.45 and in liver was 11.33 ± 0.44 while in testis 20.01 ± 1.24 within PFZ and in male muscle 19.06 ± 0.66 and liver protein content was 10.56 ± 0.51 while in testis protein content was 18.20 ± 0.64 outside PFZ during summer season from Murud fish landing center. The comparisons were made liver and gonad significant (P<0.001) and muscle to gonad (P<0.01)

whereas, the no significant value was occurred in muscle to liver within PFZ. In outside PFZ liver to gonad value was significant (P<0.001) and muscle to gonad value was similar (P<0.01) during winter season. The comparisons was made muscle to liver highly significant (P<0.05) and liver to gonad (P<0.05) and muscle to gonad significant (P<0.01) within PFZ, where as to muscle to gonad comparisons were made (P<0.01) and muscle to liver and liver to gonad value was significant in outside PFZ during summer season. Overall result showed that the protein content from winter season significantly high value than summer season, whereas, the value was highly significant within PFZ than outside PFZ. Therefore, the result showed that the protein content was increased within PFZ and decreased outside PFZ during winter and summer season.

Table 2: Protein (mg/100mg wet wt.) content in female *R. kanagurta* from Murud

Landing centre	Season	Fishing zone	Tissue		
			Muscle	Liver	Gonad
Murud	Winter	Within PFZ	21.20 ± 0.24	13.21 ± 1.02	18.20 ± 0.06 * °°
		Outside PFZ	19.05 ± 1.10	11.53 ± 0.80	18.12 ± 1.01 * °°
	Summer	Within PFZ	20.51 ± 1.11	12.42 ± 0.42	17.40 ± 0.21 *** °°
		Outside PFZ	19.16 ± 0.23	11.33 ± 0.24	16.33 ± 0.11 °°

•=P<0.001, ••= P<0.01, •••=P<0.05 comparison were made muscle to liver
 * =P<0.001, **= P<0.01, ***=P<0.05 comparison were made liver to gonad
 ° =P<0.001, °°= P<0.01, °°° =P<0.05 comparison were made muscle to gonad




Fig 2: protein (mg/100mg wet wt.) content in female *R. kanagurta* from murud

In Table. 2. & Graph. 2. Shows that in female muscle protein content was 21.20 ± 0.24 and in liver 13.21 ± 1.02 while in ovary it was 18.40 ± 0.06 within PFZ. In muscle protein content was 19.05 ± 1.10 and in liver lower to 11.53 ± 0.80 in ovary it was 18.12 ± 1.01 in outside PFZ during winter season. In muscle protein content was 20.51 ± 1.33 and in liver protein content was 12.42 ± 0.42 and in ovary it was 17.40 ± 0.21 within PFZ. From outside PFZ in muscle protein content was 19.16 ± 0.23 and in liver 11.33 ± 0.24 while in ovary it was high 16.33 ± 0.11 during summer season in Murud fish landing centre.

The comparisons were made muscle and gonad highly significant ($P < 0.05$) and muscle to liver and liver to gonad Non-significant in within PFZ. In muscle to liver highly significant value ($P < 0.05$) and liver to gonad significant ($P < 0.01$), while muscle to gonad value was high significant ($P < 0.05$) in outside PFZ during winter season. In summer season value was made muscle to gonad increased high in muscle ($P < 0.001$) highly significant and liver to gonad and muscle to liver non-significant value was observed in within PFZ. In outside PFZ value was made liver to gonad increased high in muscle ($P < 0.05$) highly significant and liver to gonad ($P < 0.05$) muscle to liver non-significant value was observed in outside PFZ.

The overall result showed that the muscle tissue of protein content was increased and p value was significantly high within PFZ than outside PFZ in winter season than summer season therefore, the result showed that the protein content was increased within PFZ and decreased outside PFZ during winter and summer season.

Discussions

The protein cycle appears to be having a strong correlation with feeding and spawning reported in a number of fish species. Maximum protein values recorded in male and female during winter season and summers season coincided with a period of intense feeding & maturing stage of this fish. This intense feeding perhaps is more in the seasons, i.e., immediately after spawning as the fish while spawning incurs energy expenditure along with the loss of gonadal elements and recoups to compensate the expenditure through vigorous feeding activity. According to Standby, (1954) [10]. also found similar observations in the trout fish. The present study showed that protein value was high within PFZ than outside PFZ among winter season than summer season. The study was observed that the seasonal variation of protein of mackerel the proportions of the components muscle tissue varied with the change of season. The

seasonal variation found in females greater than the males. Biochemical composition of any organisms are known to vary with season, size of animal, stages of maturity and availability of food, temperature etc. (Soundarapandian and Ananthan, 2008) [11]. Meanwhile, for protein contents, no significant differences were found in the *R. kanagurta* species of pelagic fish. The fish were observed to contain with mean value shows Table.1. & Fig. 1. value 11.33 to 20.40 from within PFZ while, outside PFZ value 10.56 to 19.06 In Table. 2. & Fig. 2. shows the value 11.53 to 19.05 within PFZ while in outside PFZ 11.33 to 19.16 However, the protein content could be considered as insignificant instead, as the values were derived and estimated from the difference of PFZ and outside PFZ. (Payne *et al.*, 1999; Anthony *et al.*, 2000) [12, 13].

Conclusion

The protein values calculated for different tissues based on the content of within PFZ and outside PFZ during winter and summer season from male and female. It was observed that among different tissues, protein content was maximum in the muscle of females and gonads of females, while it was minimum in the liver of both the sexes from within PFZ than outside PFZ. The present study was also revealed that within PFZ protein content was higher in males and females than the outside PFZ. The overall results showed that the protein content were increased within PFZ than outside PFZ in both sexes during winter and summer season. In conclusion, all marine fish species could be considered as a good source of protein composition and represent a very valuable essential nutrient choice for the maintenance of a healthy body of the people.

Acknowledgement

The author very thankful to the Director, Indian National Centre for Ocean Information Services (INCOIS) for financial support and an offering a gracious opportunity to work under PFZ Mission Project, Ministry of Earth Sciences, Govt. of India, Hyderabad.

References

1. Pillai VN, Satellite remote sensing applications for the benefit of coastal fisher folk – a case study. Innovations and technologies in oceanography for sustainable development, 2005, 83–84.
2. Rose DP, Connoll JM, Effects of dietary omega-3 fatty acids on human breast cancer growth and metastases in nude mice. Journal of the National Cancer Institute, 1993;85:1743–1747.

3. Marchioli R. Efficacy of n-3 polyunsaturated fatty acids after myocardial infarction Results of gissi-prevenzione Trial. *Lipids*,2001:36:119–126.
4. Marchioli R. Early protection against sudden death by n-3 polyunsaturated fatty acids after myocardial infarction: Timecourse analysis of the results of GISSI-prevenzione. *Circulation*,2002:105:1897–1903.
5. Sidhu KS, Health benefits potential risks related to consumption of fish or fish oil. *Regulatory Toxicology and Pharmacology*,2003:38:336–344.
6. Nikolsky CV, The ecology of fishes. Academic Press, 1963.
7. Lowry OH, Rosenbrough HJ, Farr AL, Randall RJ, Protein measurement with the folin phenol reagent. *Journal of Biological Chemistry*,1951:193:265–275.
8. De Zwaan A, Zandee DI, Body distribution seasonal changes in the glycogen content of the common mussel, *Mytilus edulis*. *Comparative Biochemistry and Physiology*,1972:43(0):53–58.
9. Barnes H, Blackstock J. Estimation of lipids in marine animals and tissues. Detailed investigation of the sulphophovanillin method for total lipids. *Journal of Experimental Marine Biology and Ecology*,1973:12(1):103–118.
10. Stansby ME, Composition of certain species of freshwater fish. *Food Research*,1954:16(0):231–234.
11. Soundarapandian P, Ananthan G. Effect of unilateral eyestalk ablation on the biochemical composition of commercially important juveniles of *Macrobrachium malcolmsonii*. *International Journal of Zoological Research*,2008:4(2):106–112.
12. Payne SA, Johnson BA, Otto RS, Proximate composition of some north-eastern Pacific forage fish species. *Fish Oceanography*,1999:8(3):159–177.
13. Anthony JA, Roby DD, Turco KR, Lipid content energy density of forage fishes from the Northern Gulf of Alaska. *Journal of Experimental Marine Biology and Ecology*,2000:248:53–78.